

**“Done by One” Childhood and Adolescent Immunization Schedule • New Mexico • 2011-2012**

Age ▶ Vaccine ▼	Birth	1 mo	2 mo	3 mo	4 mo	5 mo	6 mo	7 mo	8 mo	9 mo	10 mo	11 mo	12 mo (1 yr)	18 mo (1½yrs)	24 mo (2 yrs)	48 mo (4 yrs)	60 mo (5 yrs)	11-12 Yrs	13-18 Yrs
<b>DTaP (Diphtheria-Tetanus-Pertussis)</b>			Dose 1		Dose 2		Dose 3	Doses 3 and 4 should be 6 months apart				Dose 4				Dose 5			
<b>Haemophilus Influenza B (Hib)</b>			Dose 1		Dose 2		Dose 3	Dose 3 not needed if first two doses are COMVAX or Pedvax Hib. A 4 <sup>th</sup> dose may be needed for high-risk only											
<b>Hepatitis A</b>													Dose 1	Dose 2	Doses 1 and 2 should be 6 months apart				
<b>Hepatitis B</b>	At Birth		Dose 2	Dose 3 – Between 3 and 18 months															
<b>Influenza</b>	Yearly, ages 6 months to 18 years during the flu season																		
<b>Measles, Mumps &amp; Rubella</b>													Dose 1			Dose 2			
<b>Pneumonia (PCV)</b>			Dose 1		Dose 2		Dose 3	Note: First 3 doses must be 2 months apart				Dose 4							
<b>Polio</b>			Dose 1		Dose 2		Dose 3									Dose 4			
<b>Rotavirus</b>			Dose 1		Dose 2		Dose 3	If Rotarix given at 2 and 4 mos, 3 <sup>rd</sup> dose not needed											
<b>Varicella (Chicken Pox)</b>													Dose 1			Dose 2			
<b>Human Papilloma Virus (HPV)</b>																		3 Doses Female	
<b>Meningitis (Meningococcol-MCV4)</b>																		Dose 1	Dose 2
<b>Tdap (Tetanus-Diphtheria-Pertussis)</b>																		Booster	

New Mexico’s Done By One (DBO) Childhood Immunization Schedule is fully compatible with the recommendations of the Advisory Committee on Immunization Practices (ACIP) of the Centers for Disease Control and Prevention (CDC), the American Academy of Pediatrics, and the American Academy of Family Physicians. DBO simply schedules shots to be given at the earliest age allowed by the ACIP. In using this schedule, make sure that minimum intervals between shots in a series are observed. If you have any questions, call the New Mexico Immunization Hotline toll-free at 1-888-231-2367.



Shots can be given even when your child has a minor illness. If your child is not up-to-date, your child’s Primary Care Practitioner can make a plan to make sure all shots are complete. Take your child’s shot record with you to every healthcare visit.

**Your child’s Primary Care Practitioner may adjust this schedule based on what is best for your child.  
Remember, your child is not protected against these diseases until all doses have been given.**

Presbyterian exists to improve the health of the patients, members, and communities we serve.

For more information on immunizations, go to  
[www.phs.org/shots](http://www.phs.org/shots)

